

SITE:

- Lateral aspect of deltoid

- Alcohol wipe

INSERT:

- 90° angle

- Ensure no blood return

INJECT:

- Push plunger

- Remove & Massage site

DISPOSE:

- Push plunger to lock

- Place in sharps bin

ADULT

ANAPHYLAXIS	Stridor
	Bronchospasm
	Severe Abdominal Pain
	Respiratory Distress
	Tachycardia
	Shock
	Edema (Lips, Tongue, Face)
	EMT: Requires OLMC if ≥ 65 years old
	<i>Repeat after 5mins if needed</i>
RESP. DISTRESS	MEDIC ONLY
	≤ 40 years old
	Bronchospasm
	<i>OLMC for repeat</i>

6 RIGHTS OF MED. ADMINISTRATION

- Right Patient
- Right Drug
- Right Dose
- Right Time
- Right Route
- Right Documentation

PEDI

ANAPHYLAXIS	≥ 6 months old & under 55lbs (25kg)
	Skin/mucosal involvement (acute onset) with either:
	<ul style="list-style-type: none"> ● Resp. compromise ● ↓SBP
	After exposure to KNOWN patient allergen & ↓ BP
	After exposure to LIKELY allergen & 2(+) of the following:
<ul style="list-style-type: none"> ● ↓SBP ● Skin/Mucosal involvement ● GI symptoms ● Resp. Compromise 	
	<i>Repeat after 5mins if needed (OLMC required for EMTs)</i>
RESP. DISTRESS	≥ 5 years old & under 55lbs (25kg)
	Bronchospasm
	<i>Repeat after 5mins if needed</i>